



Fig. 1. Localities of records and range of *Agrionemys horsfieldii* in Iran. The range of the species in Iran and beyond is shown in grey. Own data, unless otherwise stated. – **1**- Miankaleh Peninsula 12 km W of Bandar-e-Turkmen, 36°54'N, 54°02'E, 20 m a.s.l. (KAMI 2005); **2**- Near Gorgan 36°50'N, 54°26'E (KAMI 1999); **3**- Gonbad Kavous, banks of the Atrak river (KAMI 1999); **4**- Khoshyeilagh Wildlife Refuge 36°36' to 37°02'N and 55°11' to 55°53'E, 1200 m a.s.l. (KAMI 1999); **5**- 85 km E of Shahrud, near Miyandasht, 36°25'N, 55°54'E, 1177 m a.s.l.; **6**- 115 km E of Shahrud, near Abbas Abad, 36°21'N, 56°17'E, 1270 m a.s.l.; **7**- Kalateh-rey sand dunes, 16 km NW of Zamanabad, 35°41'N, 56°38'E, 1060 m a.s.l. (CAS 228637); **8**- Sandy foothills near Sabzevar, 36°12' N, 57°30'E (KAMI 1999); **9**- East Kopetdag, 3 km NE of Mazdavand, 36°09'N, 60°35'E, 940 m a.s.l.; **10**- 13 km SW Sarakhs 36°32'N, 61°11'E (MMTT; ANDERSON 1979); **11**- Kuh-e Shakhneshin Mts, 25 km SW of Saleh Abad, 35°30'N, 60°50'E, 1690 m a.s.l.; **12**- Kuh-e Sorkh Mts, 35 km E of Dowlat Abad, 35°20'N, 59°54'E, 1830 m a.s.l.; **13**- Eshdeger range, foothills near Deyhuk, 33°15'N, 57°27'E, 1546 m a.s.l. ; **14**- Kuhistan [region centering 33°N, 57°E] (WERNER 1936); **15**- Qayen Mts, 7.2 km N of Sarbisheh, 32°39', 59°46'E, 1855 m a.s.l. (CAS 228669); **16**- Qayen Mts, Ahangarun, between Ahangarun and Doroh, 33°25'N, 60°13'E (ZARUDNOI 1903, ANDERSON 1979); **17**- Qayen Mts, near Gazik, between Ahangarun and Doroh, 32°60'N, 60°13'E (ZARUDNOI 1903); **18**- Qayen Mts, near Tabas, between Ahangarun and Doroh, 32°48'N, 60°14'E (ZARUDNOI 1903); **19**- Qayen Mts, Rud-i-Mil, between Chah-i-Sagak and Doroh, 32°17'N, 60°30'E (GRAY 1873, ANDERSON 1979); **20**- Qayen Mts, 40 km NW of Nehdandan, 31°51'N, 59°46'E., 1750 m a.s.l.; – Additional localities: Kara-iltschi Mts (NIKOLSKY 1897); Tauran (WERNER 1936); Almelh, Chodech 'Abbas, Mohammad Reza Shah National Park (MMTT).